

AD _____

AWARD NUMBER DAMD17-97-1-7242

TITLE: Identification of Protein Components of Yeast Telomerase

PRINCIPAL INVESTIGATOR: Shu-Chun Teng, Ph.D.

CONTRACTING ORGANIZATION: Princeton University
Princeton, New Jersey 08544-0036

REPORT DATE: September 1998

TYPE OF REPORT: Annual

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 4

19990617 142

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE September 1998	3. REPORT TYPE AND DATES COVERED Annual (1 Sep 97 - 31 Aug 98)	
4. TITLE AND SUBTITLE Identification of Protein Components of Yeast Telomerase			5. FUNDING NUMBERS DAMD17-97-1-7242	
6. AUTHOR(S) Shu-Chun Teng, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Princeton University Princeton, New Jersey 08544-0036			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research And Materiel Command ATTN: MCMR-RMI-S 504 Scott Street Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>Telomeres are specialized tandemly repeated DNA structures found at the natural ends of most eukaryotic chromosomes. They are required both for protecting chromosome from degradation and replicating the ends of chromosomes. Most eukaryotes replicate telomere by a special reverse transcriptase called telomerase. Another pathway of telomere formation is telomere-telomere recombination that uses homologs either in telomeric or subtelomeric repeats as the substrates. I am working on two projects involving <i>Saccharomyces cerevisiae</i> telomere formations. <i>S. cerevisiae</i> replicate their telomeres by using the telomerase pathway. In the first project I focus on understanding the nature of the <i>S. cerevisiae</i> telomerase. I found the Est2p reverse transcriptase and RNA subunit <i>TLC1</i> form a 20S telomerase complex in vivo. In the second project I tagged the subtelomere Y' element with a marker to study telomere-telomere recombination. I found that telomerase minus cells can survive through telomere-telomere recombination using either TG-rich or Y' sequences as substrates.</p>				
14. SUBJECT TERMS Breast Cancer Telomerase/Telomere/Cancer/Yeast/ <i>TLC1</i>			15. NUMBER OF PAGES 10	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

____ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

SI In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

SI In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

____ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Shu-Chun Jeng 9.3.98
PI - Signature Date

Table of contents

Front cover	1
Standard Form (SF) 298	2
Foreword	3
Table of contents	4
Introduction	5
Experimental methods	5
Assumptions, procedures, results and discussion	5
Recommendations in relation to the statement of work outlined in the proposal	9
Conclusions	10
References	10
Appendices	10

Introduction

Telomeres are specialized tandemly repeated DNA structures found at the natural ends of most eukaryotic chromosomes. They are required both for protecting chromosome from degradation and replicating the ends of chromosomes. Most eukaryotes replicate telomeres by a special reverse transcriptase called telomerase. This specialized enzyme uses a small CA-rich stretch in the RNA component as a template and the 3' hydroxyl group at chromosomal ends as a primer to extend GT-rich telomeric repeats. Another pathway of telomere formation is telomere-telomere recombination which uses homologs either in telomeric or subtelomeric repeats as substrates. In the fruit fly *Drosophila*, telomeres contain multiple retrotransposons and elongate telomeres through transpositions (Zakian, 1995).

In my original proposal I planned to use a genetic screening to look for the reverse transcriptase component of yeast telomerase, but because it was cloned right before I joined Ginger Zakian's lab (Lingner et al, 1997), I began working on two projects involving *Saccharomyces cerevisiae* telomere formations. *S. cerevisiae* replicate their telomeres by using the reverse transcriptase Est2p and RNA subunit *TLC1*, which together form a telomerase complex. In the first project I focus on understanding the nature of the *S. cerevisiae* telomerase. I will describe my findings and future directions. In the second project I tagged the subtelomere Y' element with a marker to study telomere-telomere recombination. I found that telomerase minus cells can survive through telomere-telomere recombination using either TG-rich or Y' sequences as substrates. I am writing a paper to report this finding now.

Experimental methods

All the *E.coli* and yeast operations, RT-PCR, and Southern blot analysis were performed according to "Current protocols in Molecular Biology" (Ausubel et al, 1997).

Assumptions, procedures, results and discussion

Project I. *In vitro* study of Est2p, the catalytic subunit of telomerase

To study Est2p *in vitro*, first I had to trace proteins in cell extract. I tagged the full-length EST2 with GST protein under a *GAL1* promoter in a 2-micron plasmid. Proteins were over-expressed by galactose induction and whole cell extracts were incubated with glutathione sepharose beads. After washing off unbound proteins, precipitates were subjected to RT-PCR with primers amplifying *TLC1* and *RAP1* RNAs. While no amplified RAP1 from any pull-down products and no amplified *TLC1* product from GST alone and two other proteins, CDC13p and Est1p-BP, studied in our lab, an obvious signal was amplified from GST-EST2 pull-down products (figure 1). This result suggested Est2p and *TLC1* RNA associate with each other *in vivo*.




Figure 1. RT-PCR amplification of the *TLC1* RNA from the GST-Est2p pull down.

The next question that I wanted to clarify is what is the size of telomerase. Knowing its size is important in prediction whether there are other components in the telomerase complex. To answer this question, S100 from whole cell extract were separated on a 15%-40% glycerol gradient. Fractions were collected and analyzed by RT-PCR with primers amplifying the *TLC1* RNA. PCR products were separated on a 2% agarose gel. The peak of *TLC1* was observed at the 20s position (figure 2). This result might suggest that there are other components in the telomerase complex.




Figure 2. Glycerol gradient separation of the telomerase complex.

Telomerase may form a 20s complex in vivo when it does not bind to a telomere. However, the architecture of the complex when bound to the telomere remains elusive. I have been trying to answer this question by using an in vivo chromatin immunoprecipitation assay (ChIP). This assay uses formaldehyde as a cross-linking agent, because it reacts with amino and imino groups of proteins and of DNA within minutes. After mechanic force breakage, chromatin fragments containing Est2p can be immuno-

precipitated. Cross-links can be reversed and chromatin fragments can be PCR amplified (Aparicio et al, 1997).

Using this assay, I have readily immuno-precipitated Rap1p, a telomere binding protein with telomerase. An origin binding protein in the ORC complex can also be immuno-precipitated easily and shown increased signal to bind ARS at the early S-phase (Aparicio et al, 1997). A triple HA tag right before the stop codon of the chromosomal copy of *EST2* showed a signal barely above the background. Recently I constructed an *EST2* with nine copies of Myc tag on both the chromosomal copy and a 2 micron plasmid. ChIPs of these samples are in progress.

Project II. Telomere- telomere recombination suppresses loss of telomerase

S. cerevisiae replicate their telomeres predominately by the telomerase-dependent pathway using the Est2p and RNA subunit *TLC1* to form a telomerase complex. In the *tlc1* strain, cells lose telomeres gradually and most cells die within 100 generations. However, survivors come out (Singer and Gottschling, 1994). On the Southern blots, survivors showed two distinct patterns, which is similar to Lundblad's observation in the *est1* strain (Lundblad and Blackburn, 1993). One type extends telomeres by amplifying the subtelomeric Y' elements through recombination (I called it type I). The other type contains an irregular pattern of bands. Since these bands could be hybridized by Y'-3' end probe but not other Y' regions, Lundblad and Blackburn reasoned this type of survivors is similar to type I but uses the Y'-3' end as a substrate for recombination (we called it type II) (figure 3). Given there is no obvious TG₁₋₃ sequences at the Y' 3' end between the *XhoI* site and the telomere, I assume either the Y'-3' end is not the right substrate, or that there is a novel mechanism to replicate telomeres of type II survivors.




Figure 3. Southern blot analysis of *tlc1* survivors.

What known mechanisms can generate tandem repeats of mini-Y'? The most likely possibilities are transposition and nonhomologous end joining. The structure of Y' is similar to a retrotransposon. Retrotransposons are known to transpose at *Drosophila* telomeres. Gabriel and Teng also found that retrotransposons have the ability to repair chromosomal breaks (Teng et al, 1996). I first tagged a single telomere with the *his3AI* marker and checked the histidine prototroph formation through type II recombinant formation. I observe no increase of histidine prototroph formation in *tlc1* survivors. The frequency is $\sim 1 \times 10^{-10}$ which is similar to tagging the *his3AI* marker at other place in the genome (data not shown).

Ku proteins were demonstrated to affect telomere formation (Gravel et al, 1998). To test if type II survivors replicate telomeres through nonhomologous end joining, I constructed a *tlc1 rad52* strain containing a CEN plasmid with *RAD52* and the *URA3* marker. After cells were senesced and type II survivors were recovered, the *RAD52* plasmid was lost by the FOA selection. After four re-streakings, no cell could survive.

From the previous two experiments I ruled out the possibilities of transposition and nonhomologous end joining. These results strongly suggest that telomeres of type II survivors replicate and maintain themselves through the *RAD52*-dependent homologous recombination. The substrate for homologous recombination cannot be the Y' 3' end since there is no repetitive sequences between the *XhoI* site and TG_{1,3} for homologous recombination. The substrate must be TG_{1,3} or other unknown sequences. To understand the type II recombination in detail, I traced a single telomere by studying the *his3AI* tagged telomeres. 24 type II survivors were recovered and genomic DNA were collected, digested with *XhoI* and separated on a 0.9% agarose gel. Southern blot analysis was performed with Y'-5' end, Y'-middle region, Y'-3' end, X, rDNA, LTR, C_{1,3}A and *his3AI*. As Lundblad and Blackburn reported, both Y'-3', C_{1,3}A, and *his3AI* probes can pick up multiple irregular size bands. Other probes could not hybridize these bands (data not shown).

To determine the sequence of these newly added telomeres, an inverse PCR stratagem was used (figure 4). Genomic DNAs of wild-type and two type II survivors were cut with *XhoI*. All *XhoI* digested fragments were blunted by Klenow fragment and four nucleotides. Self-ligated fragments were subjected to PCR using primer 1 and primer 2 facing opposite to each other. Expected sizes of PCR products from both wide-type and two type II survivors were obtained. These PCR products can be hybridized by C_{1,3}A probe. Sequences of the PCR products by primer 3 showed only T and G signals, whereas primer 4 showed only A and C signals on gels. Since PCR products are population heterogeneous in sequence, I tried to clone them into the pCRIITOP vector (*invitrogene*) and transformed them into four different *E. coli* strains. Surprisingly, while PCR products from wild-type telomeres were easily cloned into those *E. coli* strains, clones from telomeres of type II survivors were always difficult to get. The rarely recovered clones contained inserts between 0 and 400 bp, which are smaller than the sizes of PCR products. This result either there are repetitive sequences in the telomeres of type II survivors, or the sequence itself encodes genes toxic to bacteria. Finally I cloned

these strains into a special *E. coli* strain which can stabilize long repetitive sequences (STAB2 cells, GIBCO-BRL). The cloned plasmid showed the expected sizes. Sequences of the PCR products showed only T and G signals by primer 3 and showed only A and C signals by primer 4 on sequencing gels. Four different 4-base cutters were used to rule out the possibility that mini-Y' fragments were present in the middle of (TG₁₋₃)_n repeats. Southern blot analysis showed that while the (TG₁₋₃)_n fragment of wild-type and type I survivors can be digested into small fragments, the (TG₁₋₃)_n fragment of type II recombinants always stayed the same length (data not shown). This result suggests that the long extended telomere fragments in type II survivors are all (TG₁₋₃)_n.




Figure 4. Scheme of cloning telomeres from type II survivors.

Recommendations in relation to the statement of work outlined in the proposal.

For the first project, I intend to optimize the condition of immuno-precipitation the *EST2* strain with nine copies of Myc tag on both the chromosomal copy and a 2 micron plasmid, and to see if I can see telomerases access to telomeres in a cell-cycle regulated manner. For the second project, after submitting a paper to report the finding on telomere-telomere recombination, I plan to conduct a differential display experiment to look for genes been up-regulated in the *tlc1* strain during the survivor formation.

Conclusions

In the first project of understanding the nature of the *S. cerevisiae* telomerase. I found telomerase in vivo form a 20S complex. This complex contains both the RNA template and the reverse transcriptase. I am working on the chromatin IP technique to see if I can see telomerases access to telomeres in a cell-cycle regulated manner.

In the second project I tagged the subtelomere Y' element with a marker to study telomere-telomere recombination. I found in the telomerase knockout strain, that cells can survive through telomere-telomere recombination using either TG-rich sequence or Y' sequences as substrates. I am writing a paper to report this finding now.

References

- Aparicio et al, Components and dynamics of DNA replication complexes in *Saccharomyces cerevisiae*: redistribution of MCM proteins and Cdc45p during S phase. 1997, Cell, 91, 59-69
- Ausubel et al, Current protocols in Molecular Biology. 1997, John Wiley & Sons, Inc. press
- Gravel et al, Yeast Ku as a regulator of chromosomal DNA end structure. 1998, Science, 741-744
- Lingner et al, Reverse transcriptase motifs in the catalytic subunit of telomerase. 1997, Science 276, 561-567.
- Lundblad and Blackburn, An alternative pathway for telomere maintenance rescues est1-senescence. 1993, Cell, 73,347-360
- Singer and Gottschling, TLC1: template RNA component of *Saccharomyces cerevisiae* telomerase. 1994, Science 266, 404-409
- Teng et al, Retrotransposon reverse transcriptase-mediated repair of chromosomal breaks. 1996, Nature, 383, 641-644
- Zakian, Telomeres: beginning to understand the end. 1995, Science 270, 1601-1607.

Appendices

* not available